

# Isolation, Molecular Identification and Comparative Lipid Profiling of Microalgae and Cyanobacteria

Gopinath. S. M<sup>1</sup>, Ashalatha<sup>2</sup>, Joylin Niruba<sup>3</sup>, Meghana .R<sup>4</sup>

<sup>1</sup>Assistant Professor Department of Biotechnology, Acharya Institute of Technology, Soladevanahalli, Bangalore-560 107, Karnataka, India

<sup>2</sup>Professor & Head, Department of Biotechnology, Acharya Institute of Technology, Soladevanahalli, Bangalore-560 107, Karnataka, India

<sup>3,4</sup>Final Year B.E Biotechnology Students, Department of Biotechnology, Acharya Institute of Technology, Soladevanahalli, Bangalore-560 107, Karnataka, India

**Abstract:** *Photosynthetic micro organisms, including microalgae and cyanobacteria, are efficient in converting solar energy and recycling CO<sub>2</sub> into substrates (glycogen, triglycerides and fatty acids) for bio fuel production, and they do not compete with agriculture for arable land. As per Chisti et al [1], oil content of some microalgae exceeds 80% of the dry weight of their biomass during environmental stress such as salinity, nutritional starvation etc. Agricultural oil crops, such as soybean, oil palm and waste cooking oil are widely being used to produce biodiesel; however, oil crops produce oils in amounts that are miniscule (e.g. less than 5% of total biomass basis) compared with microalgae and cyanobacteria. In the current work, microalgae and cyanobacteria were isolated from natural sources and they purified and cultured under standard conditions. The molecular identification of the organism was done by 16s/18s rDNA technology. Lipids were extracted; particularly neutral lipids. The extracted lipids were subjected to Thin Layer Chromatography (TLC). The lipid extraction protocol was optimized for the better production of lipids which can be used as substrate for biodiesel fuel production.*

**Keywords:** Micro algae, Cyanobacteria, Bio diesel, Lipid profiling

## 1. Introduction

With a phenomenal increase in the fuel requiring processes that runs the world, the question on the sustainability of conventional fuel sources also arises. The present demand of fossil fuels reaching galactic levels has awakened a relatively new interest in alternative sources of fuel like biodiesel from microalgae. The enormous research currently happening in this field may soon open a window to a new safer world.

Many alternative sources have been identified and work is being carried on in the respective fields. Biofuel is one such. Some of the most common natural sources that can be used as feed for biofuel are seeds, corn, vegetable oils, fruits, microorganisms etc. Research has shown that biofuel from microbial sources have a higher potential than the other sources.

Biofuel, specifically biodiesel from microorganism has gained attention due to their enormous advantages over the rest. The lipid accumulating ability of certain microorganism like microalgae and cyanobacteria has made them the highest yield producing feed for biodiesel.

Microalgae are single celled, eukaryotic organisms. Owing to their oil producing nature, they are considered to be oleaginous. They produce triacylglycerides (TAGs) of favorable fatty acid chain length, an ideal feedstock for conversion to biodiesel (y. Chisti, 2007).

Cyanobacteria are aquatic eubacteria. They are photosynthetic and aquatic in nature and hence are often called "blue-green algae". This name is convenient for talking about organisms in the water that make their own

food, but does not reflect any relationship between the cyanobacteria and other organisms called algae. Cyanobacteria are relatives of the bacteria, not eukaryotes, and it is only the chloroplast in eukaryotic algae to which the cyanobacteria are related.

Both these organisms accumulate lipids in the form of free fatty acids and Tri-acyl glycerides which on transesterification with alcohols (usually methanol) produce methyl esters which is biodiesel. This biodiesel, when mixed in appropriate proportion with diesel, can be used efficiently as an environmentally safe fuel. Thus, proper considerations should be given to the drawbacks of biodiesel production as it will help in realizing the dream of producing green fuel with minimal cost and high yield making it more feasible and practically possible than it presently is.

## 2. Literature Review

Yusuf Chisti., (2007) reported that biodiesel from microalgae seems to be the only renewable biofuel that has the potential to completely displace petroleum-derived transport fuels without adversely affecting supply of food and other crop products.

Li Y et al., (2008) reported that micro algae can be potentially being employed for the production of biofuels in an economically effective and environmentally sustainable manner.

Donghui Song et al., (2008) reported the the current application of biodiesel at home and aboard and the low cost advantage of microalgae as the raw material for biodiesel is analyzed. They also reported , the role of key enzymes in

microalgal fatty acid biosynthesis and problems in manipulation of lipid production in microalgae via genetic engineering, especially proposals and measures for exploitation and utilization of oil-bearing transgenic microalgae for biodiesel production are also discussed in this review.

Laurent Lardon et al., (2009) reported the potential of microalgae as an energy source but highlighted the imperative necessity to decreasing the energy and fertilizer consumption. They also reported the nitrogen stress during culture and optimized the wet and dry extraction methods.

Tadashi Matsunga et al., (2009) reported that the strain obtained from sea water was identified using 18s rDNA technology as *Scenedesmus sp.* and it showed 73% lipid accumulation and is a promising resource that can grow as a dominant species in the sea water for the production of both liquid and solid biofuels.

Shoyan Mandal et al., (2009) reported that the presence of palmitate and oleate makes the *Scenedesmus sp.* a suitable feedstock for biodiesel production.

Luisa Gouveia et al., (2009) reported that among the various algae that they grew, *Scenedesmus obliquus* presents the most adequate fatty acid profile, namely in terms of Linolenic and other polyunsaturated fatty acids.

Li Xing et al., (2010) reported that *Scenedesmus sp.* could accumulate lipids to as high as 30% and 53% of its algal biomass in nitrogen and phosphorus starvation respectively.

Asha Parmar et al., (2011) reported that Several aspects of cyanobacteria and microalgae such as oxygenic photosynthesis, high per-acre productivity, non-food based feedstock, growth on non-productive and non-arable land, utilization of wide variety of water sources and production of valuable co-products along with biofuels have combined to capture the interest of researchers and entrepreneurs and that cyanobacteria and micro algae have a positive prospect for biofuel production.

P. Prabhakaran et al., (2011) reported different methods, including autoclaving, bead beating, microwave, sonication and a 10% NaCl solution treatments, were tested to identify the most effective cell disruption, the sonication method was found to be the most applicable and efficient method of lipid extraction from microalgae.

Bradley D et al., (2011) reported that quantitative conversion of triglycerides from several different microalgae and cyanobacteria could be achieved, including from mixed microbial biomass collected from a municipal wastewater lagoon and that these samples can be used for better biodiesel yield.

Sevgi Ertugrul Karatay et al., (2011) reported that maximum lipid contents and C16 and C18 methyl ester yields were measured as 42.8% and 46.9% for *Synechococcus sp.*, 45.0% and 67.7% for *C. aponinum*, 38.2% and 90.6% for *Phormidium sp.* The saturated compounds were 74.5%, 77.9%, 84.7% for *Synechococcus sp.*, *C. aponinum* and

*Phormidium sp.*, respectively. These crude lipids could be promising feedstock for biodiesel production.

Naira Quintana et al., (2011) reported the current state of the art in developing different types of bio energy from cyanobacteria. They also reported the major important Biochemical pathways in cyanobacteria and their possibility to influence these pathways to improve the production of specific types of energy forms including biodiesels.

Hanaa H. Abd El Baky et al., (2012) reported that by supplementing the growth of *Scenedesmus obliquus* with carbon dioxide and iron, the lipids accumulated were suitable for biodiesel production as the most abundant compounds were oleic (32.19–34.44%), Palmitic (29.54–25.12%) and Stearic (12.26–16.58% of total FAMES) acids and thus this species could be used as a suitable feedstock for biodiesel production.

Shih-Shin Ho et al., (2012) reported that under nitrogen starvation, the microalgal lipid was mainly composed of C16/C18 fatty acid (around 90%), which is suitable for biodiesel synthesis. The carbohydrate present in the biomass was mainly glucose, accounting for 77–80% of total carbohydrates. This carbohydrate composition is also suitable for fermentative biofuels production (e.g., bio ethanol and biobutanol).

Ronald Halim et al., (2012) reported that due to their high biomass productivity, rapid lipid accumulation, and ability to survive in saline water, microalgae have been identified as promising feedstocks for industrial-scale production of carbon-neutral biodiesel. They also examined the principles involved in lipid extraction from microalgal cells, a crucial downstream processing step in the production of microalgal biodiesel.

Tiara M.P et al., (2012) reported the recent publications that utilize different approaches involving engineering cyanobacteria for production of high value chemicals including biofuels.

Jordan T McEwen et al., (2012) reported that current progress of modifying microorganisms with heterogeneous elements for the production of biofuels and summarized the fatty acids production in *Escherichia coli* and cyanobacteria.

### 3. Methods / Approach

#### 3.1 Isolation and culturing conditions

The sample organisms were collected from fresh water lake in the campus of GKVK and Cuddalore district in Tamilnadu. The samples were serially diluted by taking 9.9ml of the media and 0.1ml of the sample in each tube sequentially. The dilution factor was kept constant to result in a geometric progression of concentration of the organisms in a logarithmic pattern, For each dilution, the samples were mixed thoroughly mixed before streaking on the agar plates. Once streaked, the cultures were incubated at room temperature and a light intensity of 1500lux for 15 to 20 days. This process was repeated to finally obtain one pure strain. The sample was cultured in ASM III media [2][3][4][5]. The pH was maintained at 7.3 and a light

intensity of 1500 lux was provided for their growth at room temperature. They were sub cultured every 15 days once.

### 3.2 Molecular identification-

#### 3.2.1 DNA Extraction

Cultures were pelleted by centrifuging at 10000rpm for 10 minutes. The pellet was suspended in 750 µl of suspending buffer and heated at 65°C for 45minutes. Then 650µl of lysis buffer was added and heated for 45 minutes at 65°C. This solution was then centrifuged at 10000rpm for 5minutes and the supernatant was collected in a column containing silica membrane. The columns were spun at 5X for 30-35 seconds twice. Then the membrane was washed to remove other debris and colour by using 500µl of wash buffer by spinning at 10X for 1min twice. This was followed by a dry spin at 10000rpm for 5minutes. The DNA was eluted by adding 20µl of elution buffer and heating for 2minutes followed by spinning at 10000rpm for 2minutes twice. 4µl of the collected DNA was mixed with 2µl of the gel loading dye and 1µl of the buffer and it was loaded onto the electrophoresis boat. The samples were run at 35volts for 25minutes and the concentration of the DNA obtained was noted [6].

#### 3.2.2 PCR Amplification

5µl of the extracted DNA was collected in a 1ml vials. To this 10µl of buffer, 4µl of forward primer, 4 µl of reverse primer, 4µl of dNTP's and 72µl of double distilled water was added. Primers used were 16s rDNA and 18s rDNA. The PCR machine was set for 40 cycles out of which the first 5 cycles was set for a step up process of 1°C rise for every cycle. Denaturation was set for 5 minutes at 94°C; Annealing – 94°C for 30seconds, 50°C for 30seconds, 72°C for 5seconds; for 5 min at 72°C. After setting the instrument. 1 µl of taq polymerase enzyme was added to the reaction mixture and loaded onto the PCR instrument. After the completion of the reaction, the vials were removed after the temperature of the PCR reduced to 4°C. 6µl of the sample was mixed with 2µl of the gel loading dye and 1µl of the buffer and electrophoresis was performed to check for amplified bands. Once the bands were observed, the remaining sample was run for further processing [7].

#### 3.2.3 PCR Purification

Once the bands were obtained, they were cut and treated with sodium acetate and gel elution buffer and kept for heating till the gel melted. Then 200µl of iso propanol was added and left for 5minutes. This was then take into silicon columns and spun at 5X for 1minute twice. Then the debris and colouring was removed by spinning at 10X for 1minute twice by using 500µl of wash buffer. This was followed by a dry spin at 10000rpm for 5minutes. The DNA was eluted from the silica membrane using 2µl of elution buffer and heating for 2min followed by a spin at 10000rpm for 2 minutes. 4µl of this eluted DNA was mixed with 2µl of the gel loading dye and 1µl of the buffer and run in the electrophoresis tank for 30minutes at 35volts to check for the concentration of the DNA obtained.

#### 3.2.4 Nucleotide sequence analysing using gene sequencer

The Applied Biosystems® 3130 Genetic Analyzer was used to analyze the sequence of the amplified samples in this step. This sequencer uses Sanger's Dideoxy method for

identification of the bases. In the final step, the obtained sequences were compared with Gene Bank using Basic Local Alignment Search Tool (BLAST) to identify them up to genus and species level.

#### 3.2.5 Comparison of obtained sequences with Gene Bank using BLAST

Once the chromatogram and the sequences were obtained from the gene sequencer, the sequence was copied onto the BLAST tool and the most likely species of the organism was identified and analyzed [8].

### 3.3 Lipid Extraction

The cultured organisms were subjected to three methods of lipid extraction.

#### 3.3.1 METHOD 1: General Solvent Method

Two steps were followed in this protocol:

**LYSIS:** 600OD of cells were pelleted down and 1ml of YPR buffer was added. This was incubated on ice for 20minutes. Then 1gram glass beads were added and vigorously vortexed for 30minutes. The solution was then centrifuged at 1500g for 6minutes and the lysate was separately collected. To the pellet 1ml of YPR buffer was added and vortexed for 30minutes. This was then centrifuged at 1500g for 6minutes and the lysate was collected and pooled with the lysate collected after the first spin.

**LIPID EXTRACTON:** To 5mg of lysate chloroform: methanol: acidified water (1:2:1) was added. Lipids were extracted by vortexing (keeping on ice) for 45minutes. The organic phase (bottom phase) was collected separately and 400µl of chloroform was added to the top layer and vortexed for 5 minutes. Then the organic phase was collected and they were pooled together. This phase was washed several times with acidified water till the milky interface (proteins) disappeared. The tubes were left overnight for the organic phase to evaporate.

#### 3.3.2 METHOD 2: Hexane Solvent Method

[9] 40ml broth was centrifuged and 1ml of 1M sulphuric acid was added to the pellet and kept in the water bath for 30minutes at 90°C. Then 5M sodium hydroxide was added and kept in the boiling water bath for 30minutes at 90°C. Then this was cooled and the solution was centrifuged at 10000rpm for 10minutes. The supernatant was collected and 3ml 0.5M Sulphuric acid was added to it and centrifuged at 10000rpm for 10minutes. The suspended solids were collected and 5ml hexane was added to it and kept in the boiling water bath for 15minutes at 90°C then the vials were collected and the hexane phase was collected. The vials were left overnight for the solvent to evaporate.

#### 3.3.3 METHOD 3: Bligh Dyer Method

Cells were pelleted by centrifugation at 10000rpm for 10minutes and 3.75ml of chloroform: methanol (2:1) was added to it. The solution was vortexed for 15-20 minutes. Then 1.25 ml of chloroform and 1.25 ml of distilled water was added and the top milky top layer was discarded and the bottom layer was left overnight for the solvent to evaporate [10].



### 3.4 Qualitative Analysis

Qualitative analysis was performed using Thin Layer Chromatography (TLC) [12]. The extracted lipids were dried overnight. They were then dissolved in 45µl of chloroform:methanol (2:1). 45ml of the solvent system [Petroleum ether: Diethyl ether: glacial acetic acid (70:30:1)] was used. TLC sheet of 7cm height and 4cm width was used. The standards used were linolenic acid and olive oil. 2µl of the samples and the standards were spotted onto the TLC plate and put into the glass chamber concentrated with the solvent system. The plate was left in the chamber till the solvent reached three quarters of the sheet and then the plate was transferred into the developing chamber containing iodine crystals for development and comparison of the bands.

## 4. Results / Discussion

### 4.1 Results

#### 4.1.1 MOLECULAR IDENTIFICATION

The given organism was isolated and identified at the molecular level using 16s rDNA technology as microalgae *Scenedesmus obliquus*

Forward:

TGATTCATATATTGGGTCCGTCGGGGGCTTCGTATT  
TCCATTGTCAGTAGGTGAATTCTTGGATTTATGAAA  
GTACGAACTACTGCGAAAGCATTGCCAAGGATGTT  
TTCATTAATCAAGAACGAAAGTTGGGGGCTCGAAG  
TCGATTAGATAACCGTCGTAGTCTCAACCATAAACGA  
TGCCGACTAGGGATTGGCGAATGTTTTTTAATGAC  
TTCGCCAGCACCTTATGAGAAATCAAAGTTTTTGGG  
TTCCGGGGGGAGTATGGTCGCAAGGCTGAAACTTA  
AAGGAATTGACGGAAGGGCACCACCAGGCGTGGAG  
CCTGCGGCTTAATTTGACTCAACACGGGAAAACCTTA  
CCAGGTCCAGACATAGTGAGGATTGACAGATTGAG  
AGCTCTTTCTTGATTCTATGGGTGGTGGTGCATGGC  
CGTCTTAGTTGGTGGGTTGCCTTGTCATTTGAGTAT


TTCCCGCGCAACCGGAATTTAAAGGCGGGGGGGGG  
CGGGGCTTCCGTCATCTCTTATGGACTGTCTATAT  
TTTCCGTGACTGGTCAATTTTGC GCGTGCTCACCTT  
GTGGCGCCGTCTCTCCTTCTGTAGCTTCCCCTGTTCA  
CATAACCCTTCTTGAATATCTTGACAAGATGGGGG  
TGCAGGAAGGGAGTGAGATAACCTTTCTCATAACAG  
TTGCTCATCTGTTTCGGATCGAAAGTGATCGGATAT  
TATTAAGACCGACTCACTCCTTTTTTCGTTCAACTTAA  
ATATTGAGTAAAGGCAAATACTCCTCGGAATGCTTT  
TCATTACATACTTATGTCTGCTCCTCGGATTTATTG  
GCAGCTCGCAGTGCATGGACCATTATCGGCCTTGA  
CACCGAAATG

Aligned Sequence: (628 bases)

TGATTCATATATTGGGTCCGTCGGGGGCTT  
CGTATTTCCATTGTCAGTAGGTGAATTCTT  
GGATTTATGAAAGTACGAACTACTGCGAA  
AGCATTTFCCAAGGATGTTTTTCATTAATCA  
AGAACGAAAGTTGGGGGCTCGAAGTTCGAT  
TAGATAACCGTCGTAGTCTCAACCATAAACG  
ATGCCGACTAGGGATTGGCGAATGTTTTTT  
TAATGACTTCGCCAGCACCTTATGAGAAAT  
CAAAGTTTTTGGGTTCCGGGGGGAGTATGG  
TCGCAAGGCTGAAACTTAAAGGAATTGAC  
GGAAGGGCACCACCAGGCGTGGAGCCTGC  
GGCTTAATTTGACTCAACACGGGAAAACCT  
ACCAGGTCCAGACATAGTGAGGATTGACA  
GATTGAGAGCTCTTTCTTGATTCTATGGGT  
GGTGGTGCATGGCCGTTCTTAGTTGGTGGG  
TTGCCTTGTC

The above sequence is showing maximum similarity with *Scenedesmus* sp.

The cladogram was obtained for the sp. As given below:



The given organism was isolated and identified at the molecular level using 18s rDNA technology as cyanobacteria *Synechococcus elongatus*

Forward:

CAAATTTGGATCGGGGTACAAGTTGGAAA  
CGAAGTGCTAATAACCCGATATGCCGAGA  
GGTGAATCTTTTTTGGCCTGAAGAGGAGC  
TCGCGTCTGATTAGCTAGTTGGTGGTGTAA  
GGGACCACCAAGGCGACGATCAGTAGCTG

GTCTGAGAGGAGGAGCAGCCACACTGGGA  
CTGAGACACGGCCAGACTCCTACGGGAG  
GCAGCAGTGGGGAAATTTCCGCAATGGGC  
GAAAGCCTGACGGAGCAATACCGCGTGAG  
GGAGGAAGGTCTTTGGATTGTAAACCTCTT  
TTCTCAGGGAAGAATATTGACGGTACCTGA  
GGAATAAGCACCGGCTAACTCCGTGCCAG  
CAGCCGCGGTAATACGGAGGGTGCAGCG


TTATCCGGAATTATTGGGCGTAAAGAGTCC  
 GTAGGTGGTCATGCAAGTCTGCTGTCAAAG  
 CCCACAGCTTAACTGTGGATCGGCCGGTGG  
 AAAGTGTGTGACTTGAGTACGGTAGGGGT  
 AGAGGGAATTCCCAGTGTAGCGGTGAAAT  
 GCGTAGATATTGGGAAGAACACCGGTGGC  
 GAAAGCGCTCTACTGGACCGTAACTGACA  
 CTGAGGGACGAAAGCTAAGGTAGCGAAAG  
 GGATTAGATACCCCCTTTTATTTACAGGG  
 TCTCGCTCCGTCTCCACCCGGATTTAAGA  
 GGTGCTAGTTAGTGTGTGCTCCTTATCA  
 GCAGATTTGCTGTGGACCCCTAGCGACTC  
 TTTACGCCCACTATTCCAGATATCCTGCGC  
 CCCTCCGCATTTACGCGTTGCGGCTCTGA  
 TGCAGCTGTGCTGATCTGCACGGACGTCTA  
 TAGTCTCCATGACAGATGAGCTTGACTCCA  
 ATGACTTACTCGTACGCTGCTATGATCGAC  
 AGACTAGCAGTCGTAATAACGATCGTGCG  
 TACGATAGGGATTCATGAAGAGTC

GGTGAAATCTTTTTTGGCCTGAAGAGGAGC  
 TCGCGTCTGATTAGCTAGTTGGTGGTGTA  
 GGGACCACCAAGGCGACGATCAGTAGCTG  
 GTCTGAGAGGAGGAGCAGCCACACTGGGA  
 CTGAGACACGGCCCAGACTCCTACGGGAG  
 GCAGCAGTGGGGAATTTTCCGCAATGGGC  
 GAAAGCCTGACGGAGCAATACCGCGTGAG  
 GGAGGAAGGTCTTTGGATTGTAAACCTCTT  
 TTCTCAGGGAAGAATATTGACGGTACCTGA  
 GGAATAAGCACCGGCTAACTCCGTGCCAG  
 CAGCCGCGGTAATACGGAGGGTGCAAGCG  
 TTATCCGGAATTATTGGGCGTAAAGAGTCC  
 GTAGGTGGTCATGCAAGTCTGCTGTCAAAG  
 CCCACAGCTTAACTGTGGATCGGCCGGTGG  
 AAAGTGTGTGACTTGAGTACGGTAGGGGT  
 AGAGGGAATTCCCAGTGTAGCGGTGAAAT  
 GCGTAGATATTGGGAAGAACACCGGTGGC  
 GAAAGCGCTCTACTGGACCGTAACTGACA  
 CTGAGGGACGAAAGCTAAGGTAGCGAAAG  
 GGATTAGATACCCC

Aligned Sequence: (629 bases)


CAAATTTGGATCGGGGTACAAGTTGGAAA  
 CGAAGTGCTAATAACCCGATATGCCGAGA

The above sequence is showing maximum similarity with *Synechococcus elongatus*.



4.1.2 TLC Results

The Thin Layer Chromatography was performed to qualitatively analyze the lipids extracted from the sample using respective protocols. After optimizing concentration of the standards, and the lipid extraction protocol, the lipids extracted from *Scenedesmus obliquus* and *Synechococcus elongatus* using the Bligh-Dyer method were run on TLC. Clear band separation was observed. Bands corresponding to free fatty acids and TAGs were observed by using Linolenic acid (Dilution 2) and coconut oil (Dilution 2) as standards respectively.



**Sample Preparation:** Lipid extracted using Bligh Dyer method. Sample mixed with 45 µl of chloroform: methanol (2:1)

**Lane 1:** *Scenedesmus obliquus*

**Lane 2:** *Synechococcus elongatus*

**Lane 3:** Linolenic acid (10µl of dilution 1 in 90µl of methanol- dilution 2)

**Lane 4:** coconut oil (1µl in 90 µl of methanol)

**Solvent system:** Petroleum ether: Diethyl ether: glacial acetic acid (70:30:1)

## 4.2 Discussion

The organisms are best identified at the molecular level. The sample organisms were identified and their respective phylogenetic trees were obtained which showed the evolution taking time into consideration. The mid-point shows the divergence of the species with respect to evolution. The Bligh-Dyer method was optimized as the best lipid extracting method showing clear band separation of lipids. The quantitative analysis shows the presence of clear bands for free fatty acid and TAGs in *Scenedesmus obliquus* which is in line with the recent findings of Mallick et al (2012)[13]. The TLC data clearly demonstrated the separation of the extracted lipids into different bands like Sterols, Mono acyl glycerides, Free Fatty acids, Steryl esters, Tri acyl glycerides. The neutral lipids that are useful for biodiesel production are Fatty acids and TAGs. TAGs are composed of Fatty acids. Palmitic acid (saturated) was present in maximum quantity among the different fatty acids present in *Scenedesmus* sp. But in the case of *Synechococcus* species very less free fatty acid content was observed in the TLC and Gas chromatography data which is not comparable with the findings of Atsumi et al (2013)[14]. Further medium optimization experiments were required to understand the lipid profiling of *Synechococcus elongatus*.

## 5. Conclusion

Owing to the fast depleting fossil fuel and the pollution caused by it, interest in the alternate sources of fuel has grown enormously. Biodiesel from secondary generation fuel sources like biomass has been proved to be more efficient and advantageous on many levels. However, identifying the right oleaginous species is a very step which decides the success of the fuel production and yield. Identifying the right source which can be used as feedstock for biodiesel production will bring us one step closer to making large scale production practically feasible. It also will reduce unnecessary wastage of monetary and time investment.

Comparative study of two or more species will enable narrowing down the organism which can be more suitable as feedstock. Different species have different lipid metabolism. The lipid metabolism is modified according to the environmental conditions and the related changes. Growing different species under the same conditions and using the same protocol for lipid extraction resulted in different bands in the TLC run. Free fatty acids and TAGs are used for biodiesel. The presence of clear bands corresponding to Free fatty acids and TAGs refers to better source, the quantity being proportional to the yield.

From the TLC and GC data, fatty acid content in *Scenedesmus obliquus* was found to be more than in *Synechococcus elongatus* under the given lab conditions. Palmitic acid (saturated) was present in maximum quantity among the different fatty acids present in *Scenedesmus*. Also presence of higher amount of saturated fatty acids is an advantage since they are better for biodiesel production than unsaturated fatty acids due to their lesser nitrogen emission and higher thermal efficiency. Since fatty acid content is directly proportional to the Biodiesel yield, *Scenedesmus obliquus* a more suitable organism for Biodiesel production compared to *Synechococcus elongatus* given the experimental conditions are similar.

Further study and manipulation of this species will open new possibilities for low cost large scale production in the future.

## 6. Future Scope

Micro Algae are very sensitive to changes in the environment they grow in and the lipid metabolism is in turn dependent on it. Hence, optimizing the growth conditions for better lipid accumulation will help in better yield of biodiesel. Genetic modification to increase the expression of the gene responsible for production of DAG acyl transferase enzyme which is responsible for the conversion of DAG to TAG. Perform trans-esterification of the extracted lipids (free fatty acids) to convert it into methyl esters (biodiesel). Performance test to check the Engine efficiency of the biodiesel by mixing in varying ratios with diesel. Large scale production of bio-diesel to meet the ever increasing fuel demand. Cost cutting strategies should be developed for each step of production. Lipidomic analysis of the organisms with a similar sequence to the identified species which could also be used for bio diesel production.

## Acknowledgement

I would like to thank Amnion Biosciences private limited for giving an opportunity to work in their institute and directly and indirectly helping for our work.

## References

- [1] Yusuf Chisti, Biodiesel from microalgae, Biotechnology Advances, Vol 25, May-June 2007, pp 294-306.
- [2] Mary Mennes Allen, Simple conditions for growth of unicellular blue-green algae on plates, Journal of Phycology, Vol 25, Jan 2004, pp: 1-4.
- [3] Nichols, H. W. (1973) Growth media - freshwater. In Stein, J. (Ed.) Handbook of Phycological Methods, Culture Methods and Growth Measurements, Camb. Univ. Press, Vol 24, Nov 2004 pp: 7-24.
- [4] J. B. Waterbury, J. M. Willey, Isolation and growth of marine planktonic cyanobacteria, Methods in Enzymology, Vol 167, June 1988 pp: 100-105.
- [5] Markham, J.W., Hagmeier, E, Observations on the effects of germanium dioxide on the growth of macroalgae and diatoms. Phycologia, Vol 21, Jan 1982, pp: 125-130.
- [6] Nathaniel C. Cady, Scott Stelick, Carl A. Batt, Nucleic acid purification using microfabricated silicon



structures, Biosensors and Bioelectronics, Vol 19, Oct 2003, pp 59–66.

- [7] Weisburg WG, Barns SM, Pelletier DA, Lane DJ (January 1991). "16S ribosomal DNA amplification for phylogenetic study". J Bacteriol, Vol173, July 2003 ,pp: 697–703.
- [8] Scott mcginnis, Thomas L. Madden, BLAST: at the core of a powerful and diverse set of sequence analysis tools, Nucleic Acids Research, Vol. 32, April 2004,pp 203-210.
- [9] Folch et al., A simple method for the isolation and purification of total lipids from animal tissues. J BiolChem, Vol 226, June 1957, 226, 497.
- [10] Bligh.E.G., Dyer.W.J., A rapid method for total lipid extraction and purification. Can.J.Biochem.Physiol, Vol 37, April 1959, pp:911- 917.
- [11] Ghosh A.K , N.Chauhan, S. Rajakumari, G. Daum, and R. Rajasekharan, At4g24160, aSolubleAcyl-CoenzymeA Dependent Lysophosphatidic Acid Acyltransferase, Plant Physiol, Vol 151, Oct 2009,pp: 869–881.
- [12] Feng D, Chena Z, Xuea S, Zhang W. Increased lipid production of the marine oleaginous microalgae Isochrysis Zhangiangensis (Chrsophyta) by nitrogen supplement. Bioresource Technology 2011; 102:6710-6716.
- [13] S. Mandal, S. Mallick, Biodiesel Production by the Green Microalga Scenedesmusobliquus in a Recirculatory Aquaculture System Appl Environ Microbiol, Vol 78, Aug 2012, pp: 5929–5934
- [14] McEwen JTI, Machado IM, ConnorMR, Atsumi S, Engineering Synechococcuselongatus PCC 7942 for continuous growth under diurnal conditions, Appl Environ Microbiol, Vol 78, Aug 2012, pp:1668-1675.

Prof.Ashalatha Asst Prof, Biotechnology department, Acharya Institute of Technology at Amnion Biosciences Private Limited, Bangalore. The isolation, molecular identification and lipid profiling of microalgae and cyanobacteria was carried out independently and the lipid profile was made.



**Joylin Niruba**, final year B.E. student at Department of Biotechnology, Acharya Institute of Technology, Bangalore. For the partial fulfilment of B.E. the project was carried out under the guidance of Prof. Ashalatha Asst Prof, Biotechnology department, Acharya Institute of Technology at Amnion Biosciences Private Limited, Bangalore. The isolation, molecular identification and lipid profiling of microalgae and cyanobacteria was carried out independently and the lipid profile was made.

### Author Profile



**Dr. S. M. Gopinath** received the B.Sc. and M.Sc. degrees in Microbiology from Kuvempu university, M.phil from Gulbarga university and Doctor of philosophy from Kuvempu university, Shankaraghatta, Shimoga district, Karnataka, India. As an experience in teaching for more than 2 decades and has published over 100 professional research paper in International and National refereed journals in various field of life science. He has been included as editor in various International and National journals and recognized in many professional bodies and have few DBT funded projects under him. He is now working as Professor & HOD Department of Biotechnology, Acharya Institute of Technology, Bangalore, Karnataka, India.



**Prof. Ashalatha** received the B.Sc. and M.Sc. degrees in Biotechnology from Bangalore University and pursuing Doctor of philosophy from Visvesvaraya Technological University, Belgaum, Karnataka, India. As an experience in teaching for more than 6 years and has published and presented research article in various International and National refereed journals and conferences in various field of life science. She is now working as Asst Professor and research scholar in Department of Biotechnology, Acharya Institute of Technology, Bangalore, Karnataka, India.



**Meghana R**, final year B.E. student at Department of Biotechnology, Acharya Institute of Technology, Bangalore. For the partial fulfillment of B.E. the project was carried out under the guidance of