

CBCS SCHEME

USN

--	--	--	--	--	--	--	--	--	--

15MAT31

Third Semester B.E. Degree Examination, July/August 2021 Engineering Mathematics – III

Time: 3 hrs.

Max. Marks: 80

Note: Answer any FIVE full questions.

- 1 a. Obtain the Fourier series for the function,

$$f(x) = \begin{cases} 1 + \frac{2x}{\pi} & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi} & 0 \leq x \leq \pi \end{cases}$$

Hence deduce that $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$ (08 Marks)

- b. Find the constant term and first two harmonics in the Fourier series for $f(x)$ given by the following table:

x	0	π/3	2π/3	π	4π/3	5π/3	2π
f(x)	1.0	1.4	1.9	1.7	1.5	1.2	1.0

(08 Marks)

- 2 a. Expand $f(x) = \sqrt{1 - \cos x}$ in $0 \leq x \leq 2\pi$ in a Fourier series. Evaluate $\frac{1}{1.3} + \frac{1}{3.5} + \frac{1}{5.7} + \dots$

(08 Marks)

- b. Obtain the Fourier series for $f(x) = |x|$ in $(-\ell, \ell)$ and hence evaluate $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$

(08 Marks)

- 3 a. Find the Fourier transform of $f(x) = \begin{cases} 1 - |x| & \text{for } |x| \leq 1 \\ 0 & \text{for } |x| > 1 \end{cases}$ and hence deduce that

$$\int_0^{\infty} \frac{\sin^2 t}{t^2} dt$$

(06 Marks)

- b. Find the Fourier sine transform of $e^{-|x|}$. Hence show that $\int_0^{\infty} \frac{x \sin mx}{1 + x^2} dx = \frac{\pi}{2} e^{-m}$ where

$m > 0$.

(05 Marks)

- c. Find the z-transform of (i) $(2n - 1)^2$ (ii) $\cos\left(\frac{n\pi}{2} + \frac{\pi}{4}\right)$ (05 Marks)

- 4 a. Find the Fourier transform of $f(x) = \begin{cases} 1 & |x| \leq 1 \\ 0 & |x| > 1 \end{cases}$. Hence deduce $\int_0^{\infty} \frac{\sin ax}{x} dx$. (06 Marks)

- b. Find the inverse z-transform of $\frac{4z^2 - 2z}{z^3 - 5z^2 + 8z - 4}$. (05 Marks)

- c. Solve the differential equation $u_{n+2} + 6u_{n+1} + 9u_n = 2^n$ with $u_0 = u_1 = 0$ using z-transform method. (05 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.
2. Any revealing of identification, appeal to evaluator and/or equations written eg. 42+8 = 50, will be treated as malpractice.

- 5 a. Find the coefficient of correlation and the two lines of regression for the following data:

x	1	3	4	2	5	8	9	10	13	15
y	8	6	10	8	12	16	16	10	32	32

(06 Marks)

- b. Fit a curve of the form $y = ae^{bx}$ to the following data:

x	77	100	185	239	285
y	2.4	3.4	7.0	11.1	19.6

(05 Marks)

- c. Use Regula Falsi method, find the root of the equation $x^2 - \log_e x - 12 = 0$.

(05 Marks)

- 6 a. The two regression equations of the variables x and y are $x = 19.13 - 0.87y$ and $y = 11.64 - 0.5x$. Find:

- (i) Means of x
(ii) Means of y
(iii) The correlation coefficient

(06 Marks)

- b. Fit a parabola $y = a + bx + cx^2$ to the following data:

x	-3	-2	-1	0	1	2	3
y	4.63	2.11	0.67	0.09	0.63	2.15	4.58

(05 Marks)

- c. Use Newton-Raphson method to find the real root of $3x = \cos x + 1$, take $x_0 = 0.6$ perform 2 iterations.

(05 Marks)

- 7 a. Find the cubic polynomial by using Newton forward interpolating formula which takes the following values.

x	0	1	2	3
y	1	2	1	10

(06 Marks)

- b. Apply Lagrange's formula inversely to obtain a root of the equation $f(x) = 0$ given that $f(30) = -30$, $f(34) = -13$, $f(38) = 3$, $f(42) = 18$.

(05 Marks)

- c. Use Weddle's rule to evaluate $\int_0^{\pi/2} \sqrt{\cos \theta} d\theta$ dividing the interval $\left[0, \frac{\pi}{2}\right]$ into six equal parts.

(05 Marks)

- 8 a. A survey conducted in a slum locality reveals the following interpolating information as classified below:

Income/day in rupees : x	Under 10	10-20	20-30	30-40	40-50
Number of persons : y	20	45	115	210	115

Estimate the probable number of persons in the income group 20 to 25.

(06 Marks)

- b. Using Newton divided difference formula fit an interpolating polynomial for the following data:

x	0	1	4	5
f(x)	8	11	68	123

(05 Marks)

- c. Using Simpson's $1/3^{\text{rd}}$ rule evaluate $\int_0^1 \frac{x^2}{1+x^3} dx$ taking four equal strips.

(05 Marks)

- 9 a. Find the extremal of the functional $I = \int_0^{\pi/2} (y^2 - y'^2 - 2y \sin x) dx$ under the conditions $y(0) = y\left(\frac{\pi}{2}\right) = 0$. (06 Marks)
- b. If $\vec{F} = x^2\mathbf{i} + xy\mathbf{j}$ evaluate $\int_C \vec{F} \cdot d\vec{r}$ from $(0, 0)$ to $(1, 1)$ along
 (i) the line $y = x$ (ii) the parabola $y = \sqrt{x}$ (05 Marks)
- c. Find the curve passing through the points (x_1, y_1) and (x_2, y_2) which when rotated about the x-axis gives a minimum surface area. (05 Marks)
- 10 a. Verify Green's theorem in a plane for $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$ where c is the boundary of the region enclosed by $y = \sqrt{x}$ and $y = x^2$. (06 Marks)
- b. Using divergence theorem evaluate $\int \vec{A} \cdot \hat{n} ds$ where $\vec{A} = x^3\mathbf{i} + y^3\mathbf{j} + z^3\mathbf{k}$ and s is the surface of the surface $x^2 + y^2 + z^2 = a^2$. (05 Marks)
- c. Find the geodesics on a surface given that the arc length on the surface is $s = \int_{x_1}^{x_2} \sqrt{x(1+y^2)} dx$. (05 Marks)

* * * * *