

ON $(N(k), \xi)$ -SEMI-RIEMANNIAN 3-MANIFOLDS**D.G. Prakasha**

*Department of Mathematics
Karnatak University
Dharwad – 580 003
India
e-mail: prakashadg@gmail.com*

H.G. Nagaraja

*Department of Mathematics
Central College Campus
Bangalore University
Bangalore
India
e-mail: hgnraj@yahoo.com*

G. Somashekhara

*Department of Mathematics
Acharya Institute of Technology
Soldevanahalli
Bangalore – 560107
India*

Abstract. The object of the present paper is to study 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifolds. We study $(N(k), \xi)$ -semi-Riemannian 3-manifolds which are Ricci-semi-symmetric, locally ϕ -symmetric and have η -parallel Ricci tensor.

Key words and phrases: $(N(k), \xi)$ -semi-Riemannian 3-manifold, Ricci-semi-symmetric, locally ϕ -symmetric, η -parallel Ricci tensor, η -Einstein manifold.

MSC(2000): 53C25, 53C50.

1. Introduction

Let (M, g) be an n -dimensional semi-Riemannian manifold [12] equipped with a semi-Riemannian metric g . If $\text{index}(g)=1$ then g is a Lorentzian metric and (M, g) a Lorentzian manifold [4]. If g is positive definite then g is an usual Riemannian metric and (M, g) a Riemannian manifold. The notion of $(N(k), \xi)$ -semi-Riemannian structure was introduced and studied by Tripathi and Gupta [21] to unify $N(k)$ -contact metric [3], Sasakian [5], [14], (ϵ) -Sasakian [17], [22], Kenmotsu [10], para-Sasakian [15], (ϵ) -para-Sasakian structures [20].

In this paper we study 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifolds. The paper is organized as follows. Section 2 is devoted to some basic definitions and properties of almost contact metric, almost para contact metric and $(N(k), \xi)$ -semi-Riemannian manifolds. Further, we prove that an $(N(k), \xi)$ -semi-Riemannian 3-manifold is a space form if and only if the scalar curvature r of the manifold is equal to $6k$. In Section 3, we show that a Ricci-semi-symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifold is a space-form. In Section 4, a necessary and sufficient condition for an $(N(k), \xi)$ -semi-Riemannian 3-manifold to be locally ϕ -symmetric is obtained. Section 5 contains some results on $(N(k), \xi)$ -semi-Riemannian 3-manifold with η -parallel Ricci tensor.

2. Preliminaries

Let M be an n -dimensional differentiable manifold endowed with an almost contact structure (ϕ, ξ, η) , where ϕ is a $(1, 1)$ -tensor field, ξ is a vector field and η is a 1-form on M satisfying

$$(2.1) \quad \eta(\xi) = 1, \quad \phi^2 = -I + \eta \otimes \xi,$$

where I denotes the identity transformation. It follows from (2.1) that

$$(2.2) \quad \eta \cdot \phi = 0, \quad \phi(\xi) = 0.$$

If there exists a semi-Riemannian metric g satisfying

$$(2.3) \quad g(\phi X, \phi Y) = g(X, Y) - \epsilon \eta(X)\eta(Y), \quad \forall X, Y \in \chi(M),$$

where $\epsilon = \pm 1$, then the structure (ϕ, ξ, η, g) is called an (ϵ) -almost contact metric structure and M is called an (ϵ) -almost contact metric manifold. For an (ϵ) -almost contact metric manifold, we have

$$(2.4) \quad \eta(X) = \epsilon g(X, \xi) \text{ and } \epsilon = g(\xi, \xi) \quad \forall X \in \chi(M).$$

When $\epsilon = 1$ and index of g is 0 then M is the usual Sasakian manifold and M is a Lorentz-Sasakian manifold for $\epsilon = -1$ and index of g is 1.

If $d\eta(X, Y) = g(\phi X, Y)$, then M is said to have (ϵ) -contact metric structure (ϕ, ξ, η, g) . For $\epsilon = 1$ and g Riemannian, M is the usual contact metric manifold [5]. A contact metric manifold with $\xi \in N(k)$, is called a $N(k)$ -contact metric manifold [1, 6]. If moreover, this structure is normal, that is, if

$$(2.5) \quad [\phi X, \phi Y] + \phi^2[X, Y] - \phi[X, \phi Y] - \phi[\phi X, Y] = -2d\eta(X, Y)\xi,$$

then the (ϵ) -contact metric structure is called an (ϵ) -Sasakian structure and the manifold endowed with this structure is called (ϵ) -Sasakian manifold. The physical importance of indefinite Sasakian manifolds have been pointed out by Duggal in [9].

An (ϵ) -almost contact metric structure (ϕ, ξ, η, g) is (ϵ) -Sasakian if and only if

$$(2.6) \quad (\nabla_X \phi)Y = g(X, Y)\xi - \epsilon\eta(Y)X, \quad \forall X, Y \in \chi(M),$$

where ∇ is the Levi-Civita connection with respect to g . Also we have

$$(2.7) \quad \nabla_X \xi = -\epsilon\phi X \quad \forall X \in \chi(M).$$

An almost contact metric manifold is a Kenmotsu manifold [10] if

$$(2.8) \quad (\nabla_X \phi)Y = g(\phi X, Y)\xi - \eta(Y)\phi X.$$

By (2.8), we have

$$\nabla_X \xi = X - \eta(X)\xi.$$

If in (2.1), the condition $\phi^2 = -I + \eta \otimes \xi$ is replaced by

$$(2.9) \quad \phi^2 = I - \eta \otimes \xi$$

then (M, g) is called an (ϵ) -almost paracontact metric manifold equipped with an (ϵ) -almost paracontact metric structure (ϕ, ξ, η, g) .

An (ϵ) -almost paracontact metric structure is called (ϵ) -para-Sasakian structure [20] if

$$(2.10) \quad (\nabla_X \phi)Y = -g(\phi X, \phi Y)\xi - \epsilon\eta(Y)\phi^2 X,$$

where ∇ is Levi-Civita connection with respect to the metric g . A manifold endowed with an (ϵ) -para-sasakian structure is called (ϵ) -para-Sasakian manifold [20]. For $\epsilon = 1$ and g Riemannian, M is the usual para-Sasakian manifold [15].

$(N(k), \xi)$ -semi-Riemannian manifold

The k -nullity distribution [18] of (M, g) is the distribution

$$(2.11) \quad N(k) : p \rightarrow N_p(k) = \{Z \in T_p M : R(X, Y)Z = k(g(Y, Z)X - g(X, Z)Y)\},$$

where k is a real number.

An $(N(k), \xi)$ -semi-Riemannian manifold consists of a semi-Riemannian manifold (M, g) , a k -nullity distribution $N(k)$ on (M, g) and a non-null unit vector field ξ in (M, g) belonging to $N(k)$. Through the paper we assume that $X, Y, Z, U, V, W \in \chi(M)$, where $\chi(M)$ is the Lie algebra of vector fields in M , unless specifically stated otherwise. Let ξ be a non null unit vector field in (M, g) and η its associated 1-form. Thus

$$g(\xi, \xi) = \epsilon,$$

where $\epsilon = 1$ or -1 according as ξ is spacelike or timelike, and

$$(2.12) \quad a)g(X, \xi) = \epsilon\eta(X), \quad b)\eta(\xi) = 1.$$

In an n -dimensional $(N(k), \xi)$ -semi-Riemannian manifold (M, g) , the following relations hold [21]:

$$(2.13) \quad R(X, Y)\xi = \epsilon k\{\eta(Y)X - \eta(X)Y\},$$

$$(2.14) \quad R(\xi, X)Y = \epsilon k\{\epsilon g(X, Y)\xi - \eta(Y)X\},$$

$$(2.15) \quad \eta(R(X, Y)Z) = k\{\eta(X)g(Y, Z) - \eta(Y)g(X, Z)\},$$

$$(2.16) \quad S(X, \xi) = \epsilon k(n - 1)\eta(X),$$

In a 3-dimensional Riemannian manifold we have

$$(2.17) \quad R(X, Y)Z = g(Y, Z)QX - g(X, Z)QY + S(Y, Z)X - S(X, Z)Y \\ - \frac{r}{2} [g(Y, Z)X - g(X, Z)Y],$$

where Q is the Ricci operator, i.e., $g(QX, Y) = S(X, Y)$ and r is the scalar curvature of the manifold. Putting $Z = \xi$ in (2.17) and using (2.13) and (2.16), we have

$$(2.18) \quad \epsilon(\eta(Y)QX - \eta(X)QY) = \left(-\epsilon k + \frac{r}{2}\epsilon\right) (\eta(Y)X - \eta(X)Y).$$

Putting $Y = \xi$ in (2.18) and then using (2.12(b)) and (2.16) (for $n=3$), we get

$$(2.19) \quad QX = \frac{1}{2}\{(r - 2k)X - (r - 6k)\eta(X)\xi\},$$

that is,

$$(2.20) \quad S(X, Y) = \frac{1}{2}\{(r - 2k)g(X, Y) - \epsilon(r - 6k)\eta(X)\eta(Y)\}.$$

An $(N(k), \xi)$ -semi-Riemannian manifold M is said to be η -Einstein if its Ricci tensor S is of the form

$$(2.21) \quad S(X, Y) = ag(X, Y) + b\eta(X)\eta(Y)$$

for any vector fields X, Y where a, b are functions on M . Hence from (2.20) we can state the following:

Lemma 1 *A 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifold is an η -Einstein manifold.*

By using (2.19) and (2.20) in (2.17), we obtain

$$(2.22) \quad R(X, Y)Z = \left(\frac{r}{2} - 2k\right) \{g(Y, Z)X - g(X, Z)Y\} \\ - \left(\frac{r}{2} - 3k\right) \{g(Y, Z)\eta(X)\xi - g(X, Z)\eta(Y)\xi \\ + \epsilon\eta(Y)\eta(Z)X - \epsilon\eta(X)\eta(Z)Y\}.$$

An $(N(k), \xi)$ -semi-Riemannian 3-manifold is a space of constant curvature then it is an indefinite space form.

Remark. Relations (2.19), (2.20) and (2.22) are true for

1. An $N(k)$ -contact metric 3-manifold [8] if $\epsilon = 1$,
2. A Sasakian 3-manifold if $k = 1$ and $\epsilon = 1$,
3. A Kenmotsu 3-manifold [7] if $k = -1$ and $\epsilon = 1$,
4. An (ϵ) -Sasakian 3-manifold if $k = 1$ and $\epsilon k = 1$,
5. A para-Sasakian 3-manifold [2] if $k = -1$ and $\epsilon = 1$,
6. An (ϵ) -para-Sasakian 3-manifold [19] if $k = -\epsilon$ and $\epsilon k = -1$.

Lemma 2 *A 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifold is a space form if and only if the scalar curvature $r = 6k$.*

Consequently, for a 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifold, we have the following table:

M	S =	r =
<i>$N(k)$-contact metric</i>	$\frac{1}{2}\{(r - 2k)g - (r - 6k)\eta \otimes \eta\}$	$6k$
<i>Sasakian</i>	$\frac{1}{2}\{(r - 2)g - (r - 6)\eta \otimes \eta\}$	6
<i>Kenmotsu</i>	$\frac{1}{2}\{(r + 2)g - (r + 6)\eta \otimes \eta\}$	-6
<i>(ϵ)-Sasakian</i>	$\frac{1}{2}\{(r - 2\epsilon)g - \epsilon(r - 6\epsilon)\eta \otimes \eta\}$	6ϵ
<i>para-Sasakian</i>	$\frac{1}{2}\{(r + 2)g - (r + 6)\eta \otimes \eta\}$	-6
<i>(ϵ)-para Sasakian</i>	$\frac{1}{2}\{(r + 2\epsilon)g - \epsilon(r + 6\epsilon)\eta \otimes \eta\}$	-6ϵ

Proof. Let a 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifold be an indefinite space form. Then

$$(2.23) \quad R(X, Y)Z = c\{g(Y, Z)X - g(X, Z)Y\}, \quad X, Y, Z \in \chi(M),$$

where c is the constant curvature of the manifold. By using the definition of Ricci curvature and (2.23) we have

$$(2.24) \quad S(X, Y) = 2cg(X, Y).$$

If we use (2.24) in the definition of the scalar curvature we get

$$(2.25) \quad r = 6c.$$

From (2.24) and (2.25) one can easily see that

$$(2.26) \quad S(X, Y) = \frac{r}{3}g(X, Y).$$

By plugging $X = Y = \xi$ in (2.20) and using (2.26) we obtain

$$(2.27) \quad r = 6k.$$

Conversely, if $r = 6k$, then from the equation (2.22) we can easily see that the manifold is a space form. This completes the proof. ■

3. Ricci-semi-symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifolds

A semi-Riemannian manifold M is said to be Ricci semi-symmetric [13] if its Ricci tensor S satisfies the condition

$$(3.28) \quad R(X, Y) \cdot S = 0, \quad X, Y \in \chi(M),$$

where $R(X, Y)$ acts as a derivation on S . Ricci-semisymmetric manifold is a generalization of manifold of constant curvature, Einstein manifold, Ricci symmetric manifold, symmetric manifold and semisymmetric manifold. Ricci-semisymmetric condition for Kenmotsu 3-manifolds, (ϵ) -para-Sasakian 3-manifolds and LP-Sasakian 3-manifolds are studied in [7], [19] and [16] respectively.

Let M be a Ricci-semi-symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifold. From (3.28) we have

$$(3.29) \quad S(R(X, Y)U, V) + S(U, R(X, Y)V) = 0.$$

If we put $X = \xi$ in (3.29) and use (2.14), then we get

$$(3.30) \quad kg(Y, U)S(\xi, V) - \epsilon K\eta(U)S(Y, V) + kg(Y, V)S(U, \xi) - \epsilon K\eta(V)S(U, Y) = 0.$$

By using (2.16) in (3.30) we obtain

$$(3.31) \quad \epsilon K\{2kg(Y, U)\eta(V) - \eta(U)S(Y, V) - 2kg(Y, V)\eta(U) - \eta(V)S(U, Y)\} = 0.$$

Consider that $\{e_1, e_2, e_3\}$ be an orthonormal basis of the T_pM , $p \in M$. Then, by putting $X = U = e_i$ in (2.2) and taking the summation for $1 \leq i \leq 3$, we have

$$(3.32) \quad \epsilon k\{8k\eta(V) - \epsilon S(V, \xi) - r\eta(V)\} = 0.$$

Again, by using (2.16) in (3.32), we get

$$(3.33) \quad \epsilon k(6k - r)\eta(V) = 0,$$

which gives $r = 6k$. This implies, in view of Lemma 2, that the manifold is a space form.

Therefore, we have the following:

Theorem 1 *A Ricci-semi-symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifold is a space form.*

From Theorem 1 and the above table, we can state the following corollaries:

Corollary 1 *A Ricci-semi-symmetric $N(k)$ -contact metric 3-manifold is a manifold of constant scalar curvature $6k$.*

Corollary 2 *A Ricci-semi-symmetric Sasakian 3-manifold is a manifold of constant positive scalar curvature 6.*

Corollary 3 [7] *A Ricci-semi-symmetric Kenmotsu 3-manifold is a manifold of constant negative scalar curvature -6 .*

Corollary 4 *A Ricci-semi-symmetric (ϵ) -Sasakian 3-manifold is an indefinite space form.*

Corollary 5 [2] *A Ricci-semi-symmetric para-Sasakian 3-manifold is a manifold of constant negative scalar curvature -6 .*

Corollary 6 [19] *A Ricci-semi-symmetric (ϵ) -para-Sasakian 3-manifold is an indefinite space form.*

4. Locally ϕ -symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifolds

Definition 1 An $(N(k), \xi)$ -semi-Riemannian manifold is said to be locally ϕ -symmetric if

$$\phi^2(\nabla_W R)(X, Y, Z) = 0,$$

for all vector fields W, X, Y, Z orthogonal to ξ . This notion was introduced for Sasakian manifolds by Takahashi [17].

Now, differentiating (2.22) covariantly with respect to W , we get

$$\begin{aligned} (\nabla_W R)(X, Y, Z) = & \frac{1}{2}(\nabla_W r)\{g(Y, Z)X - g(X, Z)Y - g(Y, Z)\eta(X)\xi \\ & + g(X, Z)\eta(Y)\xi - \epsilon\eta(Y)\eta(Z)X + \epsilon\eta(X)\eta(Z)Y\} \\ & - \frac{(r - 6k)}{2}\{g(Y, Z)((\nabla_W \eta)(X)\xi + \eta(X)\nabla_W \xi) \\ & - g(X, Z)((\nabla_W \eta)(Y)\xi + \eta(Y)\nabla_W \xi) \\ & + \epsilon((\nabla_W \eta)(Y)\eta(Z)X + (\nabla_W \eta)(Z)\eta(Y)X) \\ & - \epsilon((\nabla_W \eta)(X)\eta(Z)Y + (\nabla_W \eta)(Z)\eta(X)Y)\}. \end{aligned}$$

Taking W, X, Y, Z orthogonal to ξ , we have

$$\begin{aligned} (\nabla_W R)(X, Y, Z) = & \frac{1}{2}(\nabla_W r)\{g(Y, Z)X - g(X, Z)Y\} \\ (4.34) \quad & - \frac{(r - 6k)}{2}\{g(Y, Z)(\nabla_W \eta)(X)\xi - g(X, Z)(\nabla_W \eta)(Y)\xi\}. \end{aligned}$$

Applying ϕ^2 on both sides of the above equation and using $\phi \cdot \xi = 0$, we have

$$(4.35) \quad \phi^2((\nabla_W R)(X, Y, Z)) = \frac{1}{2}(\nabla_W r)\{g(Y, Z)\phi^2 X - g(X, Z)\phi^2 Y\}.$$

Now taking X, Y are orthogonal to ξ , we obtain

$$(4.36) \quad \phi^2((\nabla_W R)(X, Y, Z)) = -\frac{1}{2}(\nabla_W r)\{g(Y, Z)X - g(X, Z)Y\}$$

Hence from (4.36), we can state the following:

Theorem 2 *An $(N(k), \xi)$ -semi-Riemannian 3-manifold is locally ϕ -symmetric if and only if the scalar curvature r is constant.*

If an $(N(k), \xi)$ -semi-Riemannian 3-manifold is Ricci semi-symmetric, then we have showed that $r = 6k$, that is r is constant.

Therefore, from Theorem (2), we have

Theorem 3 *A Ricci-semi-symmetric $(N(k), \xi)$ -semi-Riemannian 3-manifold is locally ϕ -symmetric.*

5. $(N(k), \xi)$ -semi-Riemannian 3-manifold with η -parallel Ricci tensor

Definition 2 The Ricci tensor S of an $(N(k), \xi)$ -semi-Riemannian manifold M is called η -parallel if it satisfies

$$(5.37) \quad (\nabla_Z S)(\phi X, \phi Y) = 0$$

for all vector fields X, Y and Z . The notion of Ricci- η -parallelity for Sasakian manifolds was introduced by Kon in [11].

Now, let us consider a 3-dimensional $(N(k), \xi)$ -semi-Riemannian manifold with η -parallel Ricci tensor. Then, from (2.20), we get

$$(5.38) \quad S(\phi X, \phi Y) = \frac{1}{2}(r - 2k)[g(\phi X, \phi Y)].$$

Differentiating (5.38) covariantly along Z , we have

$$(5.39) \quad (\nabla_Z S)(\phi X, \phi Y) = \frac{1}{2}dr(Z)g(\phi X, \phi Y).$$

If the Ricci tensor is η -parallel, then from (5.37) and (5.39) one can get

$$\frac{1}{2}dr(Z)g(\phi X, \phi Y) = 0.$$

From which, it follows that

$$dr(Z) = 0,$$

for all Z . This leads us to the following:

Theorem 4 *Let M be an $(N(k), \xi)$ -semi-Riemannian 3-manifold with η -parallel Ricci tensor. The the scalar curvature r is constant.*

In view of Theorem (2) and Theorem (4), we have the following:

Theorem 5 *An $(N(k), \xi)$ -semi-Riemannian 3-manifold with η -parallel Ricci tensor is locally ϕ -symmetric.*

Acknowledgement. The first author (DGP) is thankful to University Grants Commission, New Delhi, India for financial support in the form of Major Research Project [F.No. 39-30/2010 (SR), dated: 23-12-2012].

References

- [1] BAGEWADI, C.S., PRAKASHA, D.G., VENKATESHA, *On pseudo projective curvature tensor of a contact metric manifold*, SUT J. Math., 43 (1)(2007), 115-126.
- [2] BAGEWADI, C.S., BASAVARAJAPPA, N.S., PRAKASHA, D.G., VENKATESHA, *On 3-dimensional para-Sasakian manifolds*, Int. e-Jour Engg. Math.: Theory and Application, 2 (2007), 110-119.
- [3] BAIKOUSSIS, CH., BLAIR, D.E., KOUFORGIORGOS, TH., *A decomposition of the curvature tensor of a contact manifold satisfying $R(X, Y)\xi = k(\eta(Y)X - \eta(X)Y)$* , Mathematical Technical Report, Univesity of Iranniana, 1992.
- [4] BEEM, J.K., EHRLICH, P.E., *Global Lorentzian geometry*, Marcel Dekker, New York, 1981.
- [5] BLAIR, D.E., *Contact manifolds in Riemannian geometry*, Lecture Notes in Mathematics, Springer-Verlag, Berlin, 509 (1976), 146.
- [6] BLAIR, D.E., KIM, J.S., TRIPATHI, M.M. *On the concircular curvature tensor of a contact metric manifold*, J. Korean Math. Soc., 42 (5)(2005), 883-892.
- [7] DE, U.C., PATHAK, G., *On 3-dimensional Kenmotsu manifolds*, Indian J. Pure Appl. Math., 35 (2)(2004), 159-165.
- [8] DE, U.C., GAZI, A.K., *On ϕ -recurrent $N(k)$ -contact metric manifolds*, Math. J. Okayama Univ., 50 (2008), 101-112.
- [9] DUGGAL, K.L., *Space time manifolds and contact structures*, Int. J. Math & Math. Sc., 13 (3) (1990), 55-553.

- [10] KENMOTSU, K., *A class of almost contact Riemannian manifold*, Tohoku Math. J., 24 (2) (1972), 93-103.
- [11] KON, M., *Invariant submanifolds in Sasakian manifolds*, Math. Annalen, 219 (1976), 277-290.
- [12] O'NEILL, B., *Semi-Rimannain geometry with applications to relativity*, Academic Press, New York, London, 1983.
- [13] MIRZOYAN, V.A., *Structure theorems for Riemannian Ric-semisymmetric spaces*, Izv. Vyssh. Uchebn. Zaved. Mat., 6 (1992), 80-89.
- [14] SASAKI, S., *On differentiable manifolds with certain structures which are closely related to almost contact structure I*, Tohoku Math. J., 12 (1960), 459-476.
- [15] SATO, I., *On a structures similar to the almost contact structure*, Tensor (N.S.), 30 (3) (1976), 219-224.
- [16] SHAIKH, A.A., DE, U.C., *On 3-dimensional LP-Sasakian manifolds*, Soochow J. Math., 26 (4) (2000), 359-368.
- [17] TAKAHASHI, T., *Sasakian manifold with pseudo-Riemannian metric*, Tohoku Math.J., 21 (1969), 271-290.
- [18] TANNO, S., *Ricci curvatures of contact Riemannian manifolds*, Tohoku Math. J., 40 (1988), 441-448.
- [19] TRIPATHI, M.M., KIHIC, E., YUKSEL PERKTAS, S., KELES, S., *On (ϵ) -para Sasakian 3-manifolds*, arXiv:0911.4786v2 [math.DG] 1 Dec 2009.
- [20] TRIPATHI, M.M., KIHIC, E., YUKSEL PERKTAS, S., KELES, S., *Indefinite almost paracontact metric manifolds*, Int. J. Math. Math Sci., doi:10.1155/2010/846195.
- [21] TRIPATHI, M.M., GUPTA, P., *On $(N(k), \xi)$ -semi-Riemannian manifolds: Semisymmetries*, Int. Electron. J. Geom., 5 (1) (2012), 42-77.
- [22] XUFENG, X., XIAOLI, C., *Two theorems on (ϵ) -Sasakian manifolds*, Int. J. Math. & Math. Sci., 21 (2) (1998), 249-254.

Accepted: 12.02.2013