



CBCS SCHEME

21MAT21

Second Semester B.E. Degree Examination, June/July 2023 Advanced Calculus and Numerical Methods

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

- 1 a. Evaluate $\int_{-c}^c \int_{-b}^b \int_{-a}^a (x + y + z) dx dy dz$. (06 Marks)
- b. Evaluate $\int_0^1 \int_x^{\sqrt{x}} xy dy dx$ by changing the order of integration. (07 Marks)
- c. Prove that $\pi(\frac{1}{2}) = \sqrt{\pi}$, using definition of Gamma function. (07 Marks)

OR

- 2 a. Evaluate $\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy$ by changing into polar coordinates. (06 Marks)
- b. Find the area between the parabolas $y^2 = 4ax$ and $x^2 = 4ay$ by using double integration. (07 Marks)
- c. Show that $\beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$. (07 Marks)

Module-2

- 3 a. Find the directional derivative of $\phi = x^2yz + 4xz^2$ at the point $(1, -2, -1)$ along $2\hat{i} - \hat{j} - 2\hat{k}$. (06 Marks)
- b. If $\vec{F} = \nabla(xy^3z^2)$, find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$ at the point $(1, -1, 1)$. (07 Marks)
- c. If $\vec{F} = (x + y + az)\hat{i} + (bx + 2y - z)\hat{j} + (x + cy + 2z)\hat{k}$, find a, b, c such that $\text{curl } \vec{F} = 0$. (07 Marks)

OR

- 4 a. If $\vec{F} = xy\hat{i} + yz\hat{j} + zx\hat{k}$, evaluate $\int_C \vec{F} \cdot d\vec{v}$ where 'c' is the curve represented by $x = t, y = t^2, z = t^3, -1 \leq t \leq 1$. (06 Marks)
- b. Using Green's theorem, evaluate $\int_C (xy + y^2) dx + x^2 dy$, where 'c' is bounded by $y = x$ and $y = x^2$. (07 Marks)
- c. Apply Stoke's theorem to evaluate $\iint \text{curl } \vec{F} \cdot \hat{n} ds$ where $\vec{F} = (x^2 + y^2)\hat{i} - 2xy\hat{j}$ taken around the rectangle bounded by the lines $x = \pm a, y = 0$ and $y = b$. (07 Marks)

Module-3

- 5 a. Form a partial differential equation by eliminating arbitrary function from $Z = f(x + at) + g(x - at)$. (06 Marks)
- b. Solve $\frac{\partial^2 z}{\partial x \partial y} = \sin x \sin y$ for which $\frac{\partial z}{\partial y} = -2 \sin y$ when $x = 0$ and $z = 0$ when y is an odd multiple of $\frac{\pi}{2}$. (07 Marks)
- c. Derive one dimensional heat equation. (07 Marks)

OR

- 6 a. Form a partial differential equation by eliminating arbitrary constant from $Z = (x - a)^2 + (y - b)^2$. (06 Marks)
- b. Solve $(y - z)p + (z - x)q = x - y$. (07 Marks)
- c. Solve $\frac{\partial^2 z}{\partial y^2} = z$ given that when $y = 0$, $z = e^x$ and $\frac{\partial z}{\partial y} = e^x$. (07 Marks)

Module-4

- 7 a. The area of a circle (A) corresponding to diameter (D) is given below:

D	80	85	90	95	100
A	5026	5674	6362	7088	7854

Find the area corresponding to diameter 105 using an appropriate interpolation formula.

(06 Marks)

- b. Find a real root of $x^3 - 2x - 5 = 0$ using Regula-Falsi method correct to 3 decimal places whose root lies between 2 and 2.5. (07 Marks)
- c. Evaluate $\int_0^{\pi/2} \sqrt{\cos \theta} d\theta$ by taking 7 ordinates by Simpson's $1/3^{\text{rd}}$ rule. (07 Marks)

OR

- 8 a. Use Newton's divided difference formula to find $f(4)$ given the data:

x	0	2	3	6
f(x)	-4	2	14	158

(06 Marks)

- b. Use Newton-Raphson method to find a real root of $x \sin x + \cos x = 0$ near $x = \pi$. Carry out the iterations upto 4 decimal places. (07 Marks)
- c. Use Lagrange's interpolation formula to find y when $x = 35$ to the following data:

x	25	30	40	60
f(x)	50	55	70	95

(07 Marks)

Module-5

- 9 a. Use the Taylor series method to find $y(0.2)$ from $\frac{dy}{dx} = y + \sin x$, $y(0) = 1$. (06 Marks)
- b. Use Runge-Kutta method of order 4, find y at $x = 0.1$, given that $\frac{dy}{dx} = 3e^x + 2y$, $y(0) = 0$ with $h = 0.1$. (07 Marks)
- c. Apply Milne's predictor-corrector method, to find $y(1.4)$ from $\frac{dy}{dx} = x^2 + \frac{y}{2}$ given that $y(1) = 2$, $y(1.1) = 2.2156$, $y(1.2) = 2.4649$, $y(1.3) = 2.7514$. (07 Marks)

OR

- 10 a. Use modified Euler's method to solve $\frac{dy}{dx} = x^2 + y$ with $y(0) = 1$, $h = 0.05$ at $x = 0.1$. (06 Marks)
- b. Use Taylor series method to find $y(0.1)$ from $\frac{dy}{dx} = x^2 + y^2$ with $y(0) = 1$. (07 Marks)
- c. Use Runge-Kutta method of 4th order, find $y(0.1)$ given that $\frac{dy}{dx} = 3x + \frac{y}{2}$, $y(0) = 1$ with $h = 0.1$. (07 Marks)
