

# CBCS SCHEME

USN

AY21ME001

21MAT21

## Second Semester B.E. Degree Examination, June/July 2023 Advanced Calculus and Numerical Methods

Time: 3 hrs.

Max. Marks: 100

*Note: Answer any FIVE full questions, choosing ONE full question from each module.*

### Module-1

- 1 a. Evaluate  $\int_{-c}^c \int_{-b}^b \int_{-a}^a (x+y+z) dx dy dz$ . (06 Marks)
- b. Evaluate  $\int_0^1 \int_x^{\sqrt{x}} xy dy dx$  by changing the order of integration. (07 Marks)
- c. Prove that  $\pi^{1/2} = \sqrt{\pi}$ , using definition of Gamma function. (07 Marks)

OR

- 2 a. Evaluate  $\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy$  by changing into polar coordinates. (06 Marks)
- b. Find the area between the parabolas  $y^2 = 4ax$  and  $x^2 = 4ay$  by using double integration. (07 Marks)
- c. Show that  $\beta(m,n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$ . (07 Marks)

### Module-2

- 3 a. Find the directional derivative of  $\phi = x^2yz + 4xz^2$  at the point  $(1, -2, -1)$  along  $2\hat{i} - \hat{j} - 2\hat{k}$ . (06 Marks)
- b. If  $\vec{F} = \nabla(xy^3z^2)$ , find  $\text{div} \vec{F}$  and  $\text{curl} \vec{F}$  at the point  $(1, -1, 1)$ . (07 Marks)
- c. If  $\vec{F} = (x+y+az)\hat{i} + (bx+2y-z)\hat{j} + (x+cy+2z)\hat{k}$ , find  $a, b, c$  such that  $\text{curl} \vec{F} = 0$ . (07 Marks)

OR

- 4 a. If  $\vec{F} = xy\hat{i} + yz\hat{j} + zx\hat{k}$ , evaluate  $\int_C \vec{F} \cdot d\vec{v}$  where 'c' is the curve represented by  $x = t, y = t^2, z = t^3, -1 \leq t \leq 1$ . (06 Marks)
- b. Using Green's theorem, evaluate  $\int_C (xy + y^2) dx + x^2 dy$ , where 'c' is bounded by  $y = x$  and  $y = x^2$ . (07 Marks)
- c. Apply Stoke's theorem to evaluate  $\iint \text{curl} \vec{F} \cdot \hat{n} ds$  where  $\vec{F} = (x^2 + y^2)\hat{i} - 2xy\hat{j}$  taken around the rectangle bounded by the lines  $x = \pm a, y = 0$  and  $y = b$ . (07 Marks)

**Module-3**

- 5 a. Form a partial differential equation by eliminating arbitrary function from  $Z = f(x + at) + g(x - at)$ . (06 Marks)
- b. Solve  $\frac{\partial^2 z}{\partial x \partial y} = \sin x \sin y$  for which  $\frac{\partial z}{\partial y} = -2 \sin y$  when  $x = 0$  and  $z = 0$  when  $y$  is an odd multiple of  $\frac{\pi}{2}$ . (07 Marks)
- c. Derive one dimensional heat equation. (07 Marks)

**OR**

- 6 a. Form a partial differential equation by eliminating arbitrary constant from  $Z = (x - a)^2 + (y - b)^2$ . (06 Marks)
- b. Solve  $(y - z)p + (z - x)q = x - y$ . (07 Marks)
- c. Solve  $\frac{\partial^2 z}{\partial y^2} = z$  given that when  $y = 0$ ,  $z = e^x$  and  $\frac{\partial z}{\partial y} = e^x$ . (07 Marks)

**Module-4**

- 7 a. The area of a circle (A) corresponding to diameter (D) is given below:

D	80	85	90	95	100
A	5026	5674	6362	7088	7854

Find the area corresponding to diameter 105 using an appropriate interpolation formula. (06 Marks)

- b. Find a real root of  $x^3 - 2x - 5 = 0$  using Regula-Falsi method correct to 3 decimal places whose root lies between 2 and 2.5. (07 Marks)
- c. Evaluate  $\int_0^{\pi/2} \sqrt{\cos \theta} d\theta$  by taking 7 ordinates by Simpson's 1/3<sup>rd</sup> rule. (07 Marks)

**OR**

- 8 a. Use Newton's divided difference formula to find  $f(4)$  given the data:

x	0	2	3	6
f(x)	-4	2	14	158

- (06 Marks)
- b. Use Newton-Raphson method to find a real root of  $x \sin x + \cos x = 0$  near  $x = \pi$ . Carry out the iterations upto 4 decimal places. (07 Marks)
- c. Use Lagrange's interpolation formula to find  $y$  when  $x = 35$  to the following data:

x	25	30	40	60
f(x)	50	55	70	95

(07 Marks)

Module-5

- 9 a. Use the Taylor series method to find  $y(0.2)$  from  $\frac{dy}{dx} = y + \sin x$ ,  $y(0) = 1$ . (06 Marks)
- b. Use Runge-Kutta method of order 4, find  $y$  at  $x = 0.1$ , given that  $\frac{dy}{dx} = 3e^x + 2y$ ,  $y(0) = 0$  with  $h = 0.1$ . (07 Marks)
- c. Apply Milne's predictor-corrector method, to find  $y(1.4)$  from  $\frac{dy}{dx} = x^2 + \frac{y}{2}$  given that  $y(1) = 2$ ,  $y(1.1) = 2.2156$ ,  $y(1.2) = 2.4649$ ,  $y(1.3) = 2.7514$ . (07 Marks)

OR

- 10 a. Use modified Euler's method to solve  $\frac{dy}{dx} = x^2 + y$  with  $y(0) = 1$ ,  $h = 0.05$  at  $x = 0.1$ . (06 Marks)
- b. Use Taylor series method to find  $y(0.1)$  from  $\frac{dy}{dx} = x^2 + y^2$  with  $y(0) = 1$ . (07 Marks)
- c. Use Runge-Kutta method of 4<sup>th</sup> order, find  $y(0.1)$  given that  $\frac{dy}{dx} = 3x + \frac{y}{2}$ ,  $y(0) = 1$  with  $h = 0.1$ . (07 Marks)

\*\*\*\*\*