



Second Semester B.E. Degree Examination, June/July 2024 Advanced Calculus and Numerical Methods

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

- 1 a. Evaluate $\int_{-1}^1 \int_0^z \int_{x-z}^{x+z} (x+y+z) dy dx dz$. (06 Marks)
- b. Evaluate $\iint_R xy dx dy$, where R is bounded by $x = 2a$, the curve $x^2 = 4ay$. (07 Marks)
- c. Prove that $\beta(m, n) = \frac{|m| |n|}{|m+n|}$ (07 Marks)

OR

- 2 a. Evaluate $\int_0^a \int_y^a \frac{x}{x^2 + y^2} dx dy$ by changing the order of integration. (06 Marks)
- b. Find the area between the parabolas $y^2 = 4ax$ and $x^2 = 4ay$. (07 Marks)
- c. Prove that $\int_0^{\frac{\pi}{2}} \sqrt{\sin \theta} d\theta \times \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{\sin \theta}} d\theta = \pi$. (07 Marks)

Module-2

- 3 a. Find the directional derivative of $Q = xy^3 + yz^2$ at the point $(2, -1, 1)$ in the direction of $i + 2j + 2k$. (06 Marks)
- b. Find $\text{div}F$ and $\text{curl}F$, where $F = \text{grad}(x^3 + y^3 + z^3 - 3xyz)$. (07 Marks)
- c. Show that $\vec{F} = \frac{x\hat{i} + y\hat{j}}{x^2 + y^2}$ is both solenoidal and irrotational. (07 Marks)

OR

- 4 a. Find the work done in moving a particle in the force field $\vec{F} = 3x^2\hat{i} + (2xz - y)\hat{j} + z\hat{k}$ along the straight line joining $(0, 0, 0)$ to $(2, 1, 3)$. (06 Marks)
- b. Apply Green's theorem to evaluate $\int_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$, where C is the boundary of the region bounded by $x = 0$, $y = 0$ and $x + y = 1$ (07 Marks)
- c. Using stoke's theorem evaluate $\int_C \vec{F} \cdot d\vec{r}$, where $\vec{F} = (x^2 + y^2)\hat{i} - 2xy\hat{j}$ and C is the boundary of the rectangle $x = \pm a$, $y = 0$, $y = b$. (07 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.
2. Any revealing of identification, appeal to evaluator and /or equations written eg. 42+8 = 50, will be treated as malpractice.

Module-3

- 5 a. Form the partial differential equation by the elimination of arbitrary functions from, $z = f(x + ay) + g(x - ay)$. (06 Marks)
- b. Derive the one dimensional heat equation. (07 Marks)
- c. Solve $(mz - ny)\frac{\partial z}{\partial x} + (nx - lz)\frac{\partial z}{\partial y} = ly - mx$. (07 Marks)

OR

- 6 a. Solve $\frac{\partial^2 z}{\partial x \partial y} = \sin x \sin y$, for which $\frac{\partial z}{\partial y} = -2 \sin y$ when $x = 0$ and $z = 0$, when y is an odd multiple of $\frac{\pi}{2}$. (06 Marks)
- b. Form the partial differential equations from $f(x + y + z, x^2 + y^2 + z^2) = 0$. (07 Marks)
- c. Solve $\frac{\partial^3 z}{\partial x^2 \partial y} + 18xy^2 + \sin(2x - y) = 0$. (07 Marks)

Module-4

- 7 a. Find a real root of the equation $x \log_{10} x = 1.2$ by regula-falsi method, correct to four decimal places. (06 Marks)
- b. Find the cubic polynomial which takes the following values by using Newton's Forward interpolation.

x	0	1	2	3
f(x)	1	2	1	10

and hence evaluate $f(4)$. (07 Marks)

- c. Evaluate $\int_0^6 \frac{dx}{1+x^2}$, by using (i) Simpson's rule (ii) Simpson's $\left(\frac{3}{8}\right)^{\text{th}}$ rule. (07 Marks)

OR

- 8 a. Using Newton's-Raphson method find real root of the equation, $3x - \cos x - 1 = 0$ near $x = 0.5$, correct to 3 decimal places. (06 Marks)
- b. Using Newton's divided difference formula for the following data :

x	5	7	11	13	17
f(x)	150	392	1452	2366	5202

Evaluate $f(9)$. (07 Marks)

- c. From the following table, estimate the number of students who obtained marks between 40 and 45.

Marks :	30 - 40	40 - 50	50 - 60	60 - 70	70 - 80
No. of students :	31	42	51	35	31

(07 Marks)

Module-5

- 9 a. Using Taylor's series method, solve $\frac{dy}{dx} = 2y + 3e^x$, find $y(0.2)$ with $y(0) = 0$ upto 4th order derivative with expansion. (06 Marks)
- b. Use Runge-Kutta method to find an approximate value of y when $x = 0.2$ given that $\frac{dy}{dx} = x + y^2$ with $y(0) = 1$ and taking $h = 0.2$. (07 Marks)
- c. Apply Milne's Predictor-corrector method, find y at $x = 0.8$ given $\frac{dy}{dx} = x - y^2$ with $y(0) = 0$, $h(0.2) = 0.02$, $y(0.4) = 0.0795$, $y(0.6) = 0.1762$ and $h = 0.2$. (07 Marks)

OR

- 10 a. Using Taylor's series method, find the value of y at $x = 0.1$ from $\frac{dy}{dx} = x^2y - 1$, $y(0) = 1$, upto 4th order derivative in the expansion. (06 Marks)
- b. Apply modified Euler's method, find $y(0.2)$ and $h = 0.2$ given $\frac{dy}{dx} = x + \sqrt{|y|}$, with $y(0) = 1$. Carry out two iterations. (07 Marks)
- c. If $\frac{dy}{dx} = 2e^x - y$, $y(0) = 2$, $y(0.1) = 2.010$, $y(0.2) = 2.04$ and $y(0.3) = 2.09$. Find $y(0.4)$ by using Milne's predictor-corrector method. (07 Marks)
