



CBCS SCHEME

BMATE201

USN

Second Semester B.E./B.Tech. Degree Supplementary Examination,
June/July 2024

Mathematics – II for EEE Stream

Time: 3 hrs.

Max. Marks: 100

- Note: 1. Answer any FIVE full questions, choosing ONE full question from each module.
2. VTU Formula Hand Book is permitted.
3. M : Marks , L: Bloom's level , C: Course outcomes.*

Module – 1			M	L	C
Q.1	a.	Find the angle between the surfaces $xy^2z = 3x + z^2$ and $3x^2 - y^2 + 2z = 1$ at the point (1, -2, 1).	7	L2	CO1
	b.	If $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$, find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$.	7	L2	CO1
	c.	Show that the vector $\vec{F} = \frac{x\hat{i} + y\hat{j}}{x^2 + y^2}$ is both solenoidal and irrotational.	6	L3	CO1
OR					
Q.2	a.	Find the total work done by the force $\vec{F} = 3xy\hat{i} - 5z\hat{j} + 10x\hat{k}$ along the curve $x = t^2 + 1$; $y = 2t^2$, $z = t^3$ from $t = 1$ to $t = 2$.	7	L2	CO1
	b.	Using Green's theorem, evaluate $\int_C (xy + y^2)dx + x^2dy$ where 'c' is the closed curve of the region bounded by $y = x$ and $y = x^2$.	7	L3	CO1
	c.	Using modern mathematical tools, write the code to find the gradient of $\phi = x^2y + 2xz - 4$.	6	L2	CO5
Module – 2					
Q.3	a.	Define a Subspace. Show that the intersection of two subspaces of a vector V is also a subspace of V.	7	L2	CO2
	b.	Show that $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(x, y, z) = (x, y, -z)$ is linear transformation.	7	L3	CO2
	c.	If $u = [2, -5, -1]^T$, $v = [-7, -4, 6]^T$, compute : i) $\langle u, v \rangle$ ii) $\ u\ ^2$ iii) $\ v\ ^2$ iv) $\ u + v\ ^2$.	6	L2	CO2
OR					
Q.4	a.	Define linearly independent and linearly dependent set of vectors. Test the vectors $v_1 = [3, 0, -6]^T$, $v_2 = [-4, 1, 7]^T$ and $v_3 = [-2, 1, 5]^T$ forms a basis.	7	L2	CO2
	b.	State Rank – Nullity Theorem. For the matrix $A = \begin{bmatrix} 1 & -4 & 9 & -7 \\ -1 & 2 & -4 & 1 \\ 5 & -6 & 10 & 7 \end{bmatrix}$, Find : i) Rank of A ii) Dim (Nul A) iii) Bases	7	L3	CO2

	c.	Using the modern mathematical tool, write the code to represent the reflection transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and to find the image of vector (10, 0) when it is reflected about y – axis.	6	L2	CO5												
Module – 3																	
Q.5	a.	Find the Laplace Transform of (i) $e^{-3t} \cos 2t$ ii) $\frac{\cos at - \cos bt}{t}$.	7	L2	CO3												
	b.	Find the Laplace Transform of the square wave function of period $2a$, defined by $f(t) = \begin{cases} k & 0 < t < a \\ -k & a < t < 2a \end{cases}$.	7	L2	CO3												
	c.	Explain $f(t) = \begin{cases} \cos t & 0 < t < \pi \\ \cos 2t & \pi \leq t < 2\pi \\ \cos 3t & t > 2\pi \end{cases}$ in term of the unit step function and hence find $L[f(t)]$.	6	L3	CO3												
OR																	
Q.6	a.	Find the inverse Laplace transformer of i) $\frac{2s-1}{s^2+4s+29}$ ii) $\frac{1}{(s-4)^2}$.	7	L2	CO3												
	b.	Using the convolution theorem, find the inverse Laplace transform of $\frac{1}{(s-1)(s^2+1)}$.	7	L3	CO3												
	c.	Solve by the Laplace transforms $y'' + k^2y = 0$, given that $y(0) = 2, y'(0) = 0$.	6	L2	CO3												
Module – 4																	
Q.7	a.	Find the real root of $x \log_{10} x = 1.2$ by Regula – Falsi method correct to 2 decimal places the root lies between (2, 3).	7	L2	CO4												
	b.	Find interpolating polynomial by Newton's divided difference formula for the data $f(1) = 4, f(3) = 32, f(4) = 55$ and $f(6) = 119$.	7	L2	CO4												
	c.	Evaluate using Simpson's $\frac{1}{3}$ rule $\int_0^6 \frac{e^x}{1+x} dx$ by taking six equal parts.	6	L2	CO4												
OR																	
Q.8	a.	Find the real root of the equation $\cos x = xe^x$, using Newton's – Raphson method, correct to 3 decimal places taking $x_0 = 0.5$.	7	L2	CO4												
	b.	Use Newton's backward interpolation formula to compute the value of y when $x = 6$, given that <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>y</td> <td>1</td> <td>-1</td> <td>1</td> <td>-1</td> <td>1</td> </tr> </table>	x	1	2	3	4	5	y	1	-1	1	-1	1	7	L3	CO4
x	1	2	3	4	5												
y	1	-1	1	-1	1												

	c.	Evaluate $\int_0^5 \frac{dx}{4x+5}$, by Trapezoidal rule, taking 6 ordinates.	6	L2	CO4
Module – 5					
Q.9	a.	Employ Taylors series method to find $y(0.2)$, given that $\frac{dy}{dx} = 2y + 3e^x$, $y(0) = 0$.	7	L3	CO4
	b.	Using Modified Euler's method, find $y(0.1)$ correct to 4 decimal places, given that $y' = x - y^2$, $y(0) = 1$, $h = 0.1$, perform 2 iterations.	7	L2	CO4
	c.	Employ Milne's predictor – corrector method given that $y' = x^2(1 + y)$ $y(1) = 1$, $y(1.1) = 1.233$, $y(1.2) = 1.548$, $y(1.3) = 1.979$ to find $y(1.4)$.	6	L3	CO4
OR					
Q.10	a.	Solve $y' = \log_{10}(x + y)$, by modified Euler's method at $x = 0.2$ and $x = 0.4$ with $h = 0.2$, perform 2 iterations at each stage.	7	L2	CO4
	b.	Use 4 th order Runge – Kutta method to solve $(x + y) y' = 1$ with $y(0.4) = 1$, at $x = 0.5$ correct to 4 decimal places.	7	L2	CO4
	c.	Using modern mathematical tools, write a code to find $y(0.1)$, given $y' = x - y$, $y(0) = 1$ by Taylors series.	6	L3	CO5
